MYCOTAXON

ISSN (print) 0093-4666 (online) 2154-8889 Mycotaxon, Ltd. ©2019

April–June 2019—Volume 134, pp. 271–273

https://doi.org/10.5248/134.271

Introduction of subfamily names for four clades in *Cladoniaceae* and *Peltigeraceae* (*Lecanoromycetes*)

H. Thorsten Lumbsch[1*] & Steven D. Leavitt[2]

[1] *Science & Education, The Field Museum, 1400 S Lakeshore Dr, Chicago, IL 60605-2496*

[2] *Department of Biology and M.L. Bean Life Science Museum, Brigham Young University, Provo, UT 84602, USA*

* Correspondence to: *tlumbsch@fieldmuseum.org*

Abstract—In a recent proposal for a classification of orders and families in *Lecanoromycetidae* and *Ostropomycetidae* based on a temporal approach, *Squamarinaceae* and *Stereocaulaceae* were synonymized with *Cladoniaceae*, and *Lobariaceae* and *Nephromataceae* with *Peltigeraceae*. Since these four synonymized families represent easily recognized, monophyletic lineages and are well-established names, we here propose to accept them at subfamilial rank, as *Lobarioideae*, *Nephromatoideae*, *Squamarinoideae*, and *Stereocauloideae*.

Key words—classification, *Lecanorales*, *Peltigerales*, phylogeny, taxonomy

Introduction

Recently we proposed a revised classification of orders and families in the two major subclasses of *Lecanoromycetes*, *Lecanoromycetidae* and *Ostropomycetidae* (Kraichak & al. 2018) based on a temporal banding approach (Avise & Johns 1999, Kraichak & al. 2017). This classification proposal differed at the family level from other recently proposed classifications (Jaklitsch & al. 2016; Lücking & al. 2016, 2017; Lumbsch & Huhndorf 2010; Wijayawardene & al. 2018) in accepting some clades at family level (such as *Diploschistaceae*, *Thelotremataceae*, and *Varicellariaceae*) while synonymizing other clades previously accepted as families within older families. The latter clades represented monophyletic lineages but were shown to have diverged more recently than 111–235 Mya,

which was the temporal band identified for the family rank. Four of the families synonymized in Kraichak & al. (2018) have been widely used names, and, hence, we suggest they should be accepted at subfamily rank. This allows recognition of these well-established and monophyletic clades while avoiding confusion due to incomparable ranks of clades.

Below we formally propose new subfamilies for the previously accepted families *Lobariaceae*, *Nephromataceae*, *Squamarinaceae*, and *Stereocaulaceae*. With the description of these subfamilies, *Cladonioideae* will become an available name for *Cladoniaceae* s.str. (i.e., excluding *Squamarinoideae* and *Stereocauloideae*) and *Peltigeroideae* for *Peltigeraceae* s.str. (i.e., excluding *Lobarioideae* and *Nephromatoideae*).

Taxonomy

Cladoniaceae Zenker, in Goebel & Kunze, Pharmac. Waarenk. 1(3): 124 (1827).

Squamarinoideae Lumbsch & S.D. Leav., **stat. nov.**
MB 830561
Basionym: *Squamarinaceae* Hafellner, Beih. Nova Hedwigia 79: 342 (1984).

Stereocauloideae Lumbsch & S.D. Leav., **stat. nov.**
MB 830562
Basionym: *Stereocaulaceae* Chevall., Fl. Gén. Env. Paris 1: 596 (1826).

Peltigeraceae Dumort., Comment. Bot.: 68 (1822).

Lobarioideae Lumbsch & S.D. Leav., **stat. nov.**
MB 830563
Basionym: *Lobariaceae* Chevall., Fl. Gén. Env. Paris 1: 609 (1826).

Nephromatoideae Lumbsch & S.D. Leav., **stat. nov.**
MB 830564
Basionym: *Nephromataceae* Wetmore ex J.C. David & D. Hawksw., Syst. Ascom. 10: 15 (1991).

Acknowledgements

We wish to thank Dr. Brendan Hodkinson (Janssen Research & Development, Spring House, PA USA) and Dr. Christian Printzen (Senckenberg Research Institute, Frankfurt/Main, Germany) for reviewing this manuscript.

Literature cited

Avise JC, Johns GC. 1999. Proposal for a standardized temporal scheme of biological classification for extant species. Proceedings of the National Academy of Science of the United States of America 96: 7358–7363. https://doi.org/10.1073/pnas.96.13.7358

Jaklitsch WM, Baral HO, Lücking R, Lumbsch HT. 2016. Syllabus of plant families — A. Engler's Syllabus der Pflanzenfamilien, Part 1/2 *Ascomycota*. Gebr. Borntraeger Verlagsbuchhandlung. 322 p.

Kraichak E, Crespo A, Divakar PK, Leavitt SD, Lumbsch HT. 2017. A temporal banding approach for consistent taxonomic ranking above the species level. Scientific Reports 7:2297 [7 p.]. https://doi.org/10.1038/s41598-017-02477-7

Kraichak E, Huang JP, Nelsen M, Leavitt SD, Lumbsch HT. 2018. A revised classification of orders and families in the two major subclasses of *Lecanoromycetes* (*Ascomycota*) based on a temporal approach. Botanical Journal of the Linnean Society 188: 233–249. https://doi.org/10.1093/botlinnean/boy060

Lücking R, Hodkinson BP, Leavitt SD. 2016. The 2016 classification of lichenized fungi in the *Ascomycota* and *Basidiomycota* – approaching one thousand genera. Bryologist 119: 361–416. https://doi.org/10.1639/0007-2745-119.4.361

Lücking R, Hodkinson BP, Leavitt SD. 2017. Corrections and amendments to the 2016 classification of lichenized fungi in the *Ascomycota* and *Basidiomycota*. Bryologist 120: 58–69. https://doi.org/10.1639/0007-2745-120.1.058

Lumbsch HT, Huhndorf SM. 2010. Myconet volume 14. Part one. Outline of *Ascomycota*—2009. Fieldiana Life and Earth Sciences 1: 1–42. https://doi.org/10.3158/1557.1

Wijayawardene NN, Hyde KD, Lumbsch HT, Liu JK, Maharachchikumbura SSN, Ekanayaka AH, Tian Q, Phookamsak R. 2018. Outline of *Ascomycota*: 2017. Fungal Diversity 88: 167–263. https://doi.org/10.1007/s13225-018-0394-8